

**12th Conference on
Retroviruses and Opportunistic
Infections (CROI)
Boston MA, USA
February 22 – 25, 2005**

**COMPARISON OF HIV DECAY
RATES WITH THE ADDITION OF
ENFUVIRTIDE TO A
COMBINATION OF BOOSTED
SAQUINAVIR AND EFAVIRENZ IN
PI- AND NNRTI-NAIVE PATIENTS:
A RANDOMIZED, CONTROLLED
TRIAL**

***Boyd MA^{1,2}, Perelson AS³, Dixit NM³,
Siangphoe U¹, Duncombe CJ^{1,2}, Burger D⁴,
Ubolyam S¹, Mahanontharit A¹, Paksin S¹,
Phanuphak P⁵, Lange JMA^{1,6},
Cooper DA^{1,2} and Ruxrungtham K^{1,5}***

¹ HIV-NAT, TRCARC, Bangkok, Thailand;

*² NCHECR, University of New South Wales,
Sydney, Australia;*

³ Los Alamos National Laboratory, New Mexico, USA;

*⁴ Nijmegen University Centre, University Medical
Centre Nijmegen, the Netherlands;*

*⁵ Faculty of Medicine, Chulalongkorn University,
Bangkok, Thailand;*

⁶ IATEC and AMC, Amsterdam, the Netherlands